

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Journal of Diagnostic Pathology and Oncology

Journal homepage: <https://www.jdpo.org/>

Case Report

Juvenile fibroadenoma of the breast

Sunil Vitthalrao Jagtap^{1,*}, Swati Sunil Jagtap², Nidhi Rajubhai Goswami¹,
Neha Babasaheb Desai¹

¹Dept. of Pathology, Krishna Institute of Medical Sciences, Karad, Maharashtra, India

²Dept. of Physiology, Krishna Institute of Medical Sciences, Karad, Maharashtra, India



ARTICLE INFO

Article history:

Received 25-07-2022

Accepted 08-08-2022

Available online 22-09-2022

Keywords:

Benign breast tumors

Juvenile fibroadenoma

Mastectomy

ABSTRACT

Juvenile fibroadenoma is a rare variant of fibroadenoma and is characterized by rapidly enlarging, painless, and unilateral masses. We presents a case of juvenile fibroadenoma of the breast in a 14 year old girl, presented with rapidly growing mass in the right breast of four months duration. On local examination a solitary, firm, mobile, unilateral, non-tender mass was felt in the upper and lower outer quadrant of the right breast, measuring 6.5 x 4.1 x 3.5cm. The overlying skin was normal. There was no axillary lymphadenopathy. The sonomammography of the right breast showed a single, oval, hypoechoic, well circumscribed mass lesion with cleft like spaces features suggestive of complex fibroadenoma BIRADS IV a. On fine niddle aspiration cytology showed sheets of hyperplastic benign ductal epithelial cells with myoepithelial cells and a background of benign bipolar nuclei and blood, without inflammatory cells. It was suggestive of fibroadenoma. She underwent right lumpectomy with breast conservation. We received right breast lumpectomy specimen. On gross showed a well-circumscribed, grey white, firm mass m 6.3 x 4.1 x 3.2 cm. On histologic features reported as juvenile fibroadenoma right breast. The postoperative period was uneventful and she recovered well and advise regular follow up.

Conclusion: Juvenile fibroadenomas are rare benign breast tumors that should be distinguished from other masses of breast. Herewith we are presenting case of Juvenile fibroadenoma for its clinical, radio imaging, pathological features and management.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Breast masses in children and adolescents are rare.¹ The fibroadenoma being the most frequent breast tumor in adolescent girls. Fibroadenoma is termed juvenile if it occurs in children and adolescents between the ages of 10–18 years.² Juvenile fibroadenoma rapidly grows to a large size and the patient presents with breast enlargement, nipple displacement, overlying skin stretching, pain etc. In few cases they shows a prominent vasculature. On ultrasonography it is usually solitary, homogeneous, and

hypoechoic mass. They can result in an esthetic problem due to breast asymmetry or hypertrophy.³ In the majority of pediatric breast lesions they are of benign etiology, therefore the management approach should be proper.

2. Case Report

A 14-year-old female patient presented to our surgery department having a rapidly growing mass in the right breast in the upper outer quadrant since last four month. There was no any history of trauma, fever, nipple discharge, anorexia. There was no significant family history. No history of any hormonal intake or irradiation to the chest. She has regular menstruation since last one year.

* Corresponding author.

E-mail address: drsvjagtap@gmail.com (S. V. Jagtap).




Fig. 1: Sonomammography right breast showed a 6 x 6 x 5cm, oval, hypoechoic, well circumscribed mass lesion with cleft like spaces.



Fig. 2: Cut section right breast mass well circumscribed, grey white with slit like spaces.

She noted right breast mass associated with pain during menstruation. On local examination, a firm, mobile, solitary and unilateral, non-tender mass was felt in the upper outer quadrant of the right breast, measuring 6.5 x 4.1 x 3.5cm. The overlying skin was normal. There was no axillary lymphadenopathy. The left breast was normal. Routine hematological and biochemical parameters were within normal limits. The sonomammography of the right breast showed a 6 x 6 x 5 cm, oval, hypoechoic, well circumscribed mass lesion with cleft like spaces. Features suggestive of complex fibroadenoma -BIRADS IVa. (Figure 1). On fine needle aspiration cytology showed sheets of hyperplastic




Fig. 3: showing increased epithelial, stromal cellularity with pericanalicular growth pattern.(H& E stain,40x)




Fig. 4: Breast tissue showing mild epithelial hyperplasia and increased stromal hypercellularity. (H& E stain,100x)

benign ductal epithelial cells with myoepithelial cells and a background of benign bipolar nuclei and blood, without inflammatory cells. It was reported as benign fibroepithelial tumor suggestive of fibroadenoma. She underwent right lumpectomy with breast conservation. The postoperative period was uneventful, and she recovered well and advise regular follow up.

We received right breast lumpectomy specimen. On gross examination a firm, solitary, circumscribed, mass of the right breast measuring 6.3 x 4.1 x 3.5 cm was noted. On cut section grey white, firm, with cleft like spaces was noted (Figure 2). There was no areas of any necrosis or hemorrhages.

On microscopy breast tissue showing well-circumscribed lesion having uniform biphasic stromal and epithelial proliferation with predominantly pericanalicular growth patterns. Glandular elements have intact myoepithelial cell layer. An areas hypercellular stroma was noted, in focal areas epithelial hyperplasia, apocrine change was noted. The leaf-like fronds are not seen. There was no nuclear atypia. On histologic features reported as juvenile fibroadenoma right breast. (Figures 3 and 4).

3. Discussion

In adolescent girls the breast masses are rare and mostly are of benign natures. Juvenile fibroadenomas account for 0.5% and 4% of all fibroadenomas.⁴ Juvenile fibroadenoma is a rare variant of fibroadenoma and is characterized by rapidly enlarging, painless, and unilateral masses occurring at ages between 10 and 18 years. Juvenile fibroadenoma shows rapid growth and large size as noted in our case in 14 year girl having the duration was 4 months with large breast mass of size of 6.3 cm.

These are mainly unilateral, but rarely bilateral juvenile fibroadenomas are reported.⁵ The etiology of the disease is still not well understood, but hormonal factors have been proposed.

Clinically they are characterized by massive and rapid enlargement of breast. The fibroadenoma of breast have variants like cellular, juvenile, myxoid, complex fibroadenoma etc. The Juvenile fibroadenoma shows increased epithelial hyperplasia rearely with micropapillary projections. There is increased stromal cellularity, namely pericanalicular growth pattern with fascicular arrangement. These all microscopic findings were noted in our case. The incidence of malignant transformation ranges from 0.002% to 0.1% cases.⁶

Giant juvenile fibroadenomas are found in in 0.5% of all fibroadenomas.⁷ Giant fibroadenoma tumors are considered when >500 g or disproportionally large compared to rest of breast. Rarely multiple juvenile fibroadenomas are noted.⁸

Fibroadenomas with hypercellular stroma and prominent intracanalicular pattern can show morphologic overlap with benign phyllodes tumors. The diffrential diagnosis

are phyllodes tumor, virginal hypertrophy, juvenile papillomatosis, etc.

For the diagnosis on the FNAC breast about 97.4% accuracy in lesion of epithelial proliferative lesion without atypia was noted, however the histopathology is the gold standard for diagnosis.⁹ The phyllodes tumor of breast differs from giant fibroadenoma by the presence of leaf-like structures and stromal cell atypia on histopathological evaluation .Prominent mitotic activity $\geq 3/10$ high power fields are noted. The tumor may infiltrate surrounding breast. In the virginal hypertrophy the growth is diffuse, not circumscribed and on microcopy it is characterized by abundant connective tissue with duct proliferation and lack of lobule formation. In cases of juvenile papillomatosis stroma is paucicellular, along with many time there are prominent cysts formation is noted.

Management of juvenile fibroadenoma ranges from tumor excision to simply observation, since complete tumor regression may occur in 10-59% of lesions.¹⁰ The treatment is complete excision of the tumor, preservation of the areola and nipple, and achievement of symmetrical breasts. Surgical intervention is indicated for fibroadenomas that are more than 5 cm in diameter, rapidly growing, cause severe pain, distort the breast architecture, or lead to overlying skin changes. Clinical assessment followed by histopathological examination provides an accurate diagnosis of breast fibroadenomas.¹¹

4. Conclusion

Juvenile fibroadenomas are rare benign breast tumors that should be distinguished from other breast neoplasms. Herewith we are presenting case of Juvenile fibroadenomas for its clinical, radioimaging, pathological features and management. Juvenile fibroadenomas may be management challenge in adolescent.

5. Conflict of Interest

The authors declare no relevant conflicts of interest.

6. Source of Funding

None.


References


1. Kaneda HJ, Mack J, Kasales CJ, Schetter S. Pediatric and adolescent breast masses: a review of pathophysiology, imaging, diagnosis, and treatment. *AJR Am J Roentgenol*. 2013;200(2):204–12. doi:10.2214/AJR.12.9560.
2. Wechselberger G, Schoeller T, Piza-Katzer H. Juvenile fibroadenoma of the breast. *Surgery*. 2002;132(1):106–7. doi:10.1067/msy.2002.117375.
3. Cerrato F, Labow BI. Diagnosis and management of fibroadenomas in the adolescent breast. *Semin Plast Surg*. 2013;27(1):23–5. doi:10.1055/s-0033-1343992.
4. Ng WK, Mrad MA, Brown MH. Juvenile fibroadenoma of the breast: treatment and literature review. *Can J Plast Surg*. 2011;19(3):105–7.


- doi:10.1177/229255031101900308.
5. Mukhopadhyay M, Patra R, Mondal S, Ghosh A, Ray AK. Bilateral giant juvenile fibroadenoma of breasts. *J Indian Assoc Pediatr Surg.* 2009;14(2):68–9. doi:10.4103/0971-9261.55156.
 6. Wu YT, Chen ST, Chen CJ, Kuo YL, Tsenq LM, Chen DR, et al. Breast cancer arising within fibroadenoma: collective analysis of case reports in the literature and hints on treatment policy. *World J Surg Oncol.* 2014;12:335. doi:10.1186/1477-7819-12-335.
 7. Baxi M, Agarwal A, Mishra A, Agarwal G, Mishra SK. Multiple Bilateral Giant Juvenile Fibroadenomas of Breast. *Eur J Surg.* 2000;166(10):828–30. doi:10.1080/110241500447515.
 8. Samala DS, Gedam M. Multiple fibroadenomas in single breast. *J Case Rep.* 2015;5(1):140–2.
 9. Jagtap SV, Beniwal A, Shah HP, Jagtap SS. Application of Probabilistic Approach in Reporting Breast Lesions on Fine Needle Aspiration Cytology: Institutional Study. *National J Lab Med.* 2016;5(2):17–21. doi:10.7860/NJLM/2016/17521:2105.
 10. Jayasinghe Y, Simmons PS. Fibroadenomas in adolescence. *Curr Opin Obstet Gynecol.* 2009;21(5):402–6. doi:10.1097/GCO.0b013e32832fa06b.
 11. Boral S, Jagtap SV. Clinicohistopathological study of benign breast lesions in surgically excised specimens in a tertiary care hospital. *J Can Res Ther.* 2021;doi:10.4103/jcrt.JCRT_688_20.

Author biography

Sunil Vitthalrao Jagtap, Professor  <https://orcid.org/0000-0002-4844-7624>

Swati Sunil Jagtap, Associate Professor  <https://orcid.org/0000-0001-8905-6163>

Nidhi Rajubhai Goswami, Assistant Lecturer  <https://orcid.org/0000-0002-4342-4028>

Neha Babasaheb Desai, Assistant Lecturer  <https://orcid.org/0000-0001-6089-4323>

<p>Cite this article: Jagtap SV, Jagtap SS, Goswami NR, Desai NB. Juvenile fibroadenoma of the breast. <i>IP J Diagn Pathol Oncol</i> 2022;7(3):183-186.</p>
