

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Journal of Diagnostic Pathology and Oncology

Journal homepage: <https://www.jdpo.org/>

Case Report

Acquired lymphangioma circumscriptum of the vulva in a middle aged woman with chronic hidradenitis suppurativa – A case report

Neelam Gupta¹, Preeti Joseph^{1,*}, Lav Behl¹, Mehak¹, Sanjeev Uppal²

¹Dept. of Pathology, Maharishi Markandeshwar Medical College and Hospital, Solan, Himachal Pradesh, India

²Dayanand Medical College & Hospital, Ludhiana, Punjab, India



ARTICLE INFO

Article history:

Received 23-01-2023

Accepted 20-02-2023

Available online 29-03-2023

Keywords:

Hidradenitis suppurativa

Lymphangioma circumscriptum

Vulva

ABSTRACT

Lymphangioma circumscriptum (LC) is a distention of lymphatic channels present in skin and subcutaneous tissues. LC is either primary (usually present at birth or develops in early childhood) or secondary due to diminished lymphatic outflow. It is commonly prone to develop in oral cavity, tongue, proximal regions of arms and legs, groin, axilla, and trunk. Primary vulvar involvement is unusual. Vulvar LC is a rare untoward outcome of hidradenitis suppurativa (HS), which causes significant physical and emotional discomfort. We report a case of acquired LC of vulva as a sequelae of HS in a 42 years old female who presented with painful papules, edema and thickening of the vulva with history of foul-smelling discharging sinuses. The patient was managed surgically by wide excision and reconstructive surgery by Plastic surgery unit.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Acquired lymphangioma or lymphangiectasis rarely involve vulva.¹ Most commonly acquired genital lymphedema occurs due to radiation or surgical treatment for cervical cancer, infective causes (acute cellulitis, filariasis, and genital tuberculosis). Lymphedema in context to chronic Hidradenitis suppurativa is infrequent.^{1,2} HS is a chronic inflammatory disorder characterized by recurrent abscesses, draining sinus tracts, scarring, and blockage of lymphatic channels and accumulation of lymph in the interstitial space. The most characteristic clinical presentation is of multiple small raised bleb-like lesions from which watery fluid may ooze. A 42 years old female presented in skin OPD with a history of painful papular lesions, edema and foul smelling pus discharging sinuses of vulvar region. The patient was treated surgically with wide excision and skin grafting.

* Corresponding author.

E-mail addresses: neelamgupta353@gmail.com (P. Joseph), <https://orcid.org/0000-0002-0290-6338> (Mehak).

2. Case Report

A 42-year-old female presented in skin OPD with complaint of painful erythematous swelling with thickened skin and foul-smelling discharging sinuses, over vulvar region for the past one year (Figure 1). There was no history of swelling in lower limb, any abdominal surgery or any intrauterine device, pulmonary tuberculosis or diabetes. Patient was referred to Plastic Surgery unit where she underwent reconstructive surgery with local wide excision of edematous vulvar skin followed by skin grafting. Grossly, skin covered soft tissue received in histopathology section measuring 23x10x9cm. The external surface showed multiple pus discharging sinuses along with pustules (Figure 2). Cut section showed solid and partially cystic areas containing purulent material.

Microscopically, epidermis showed pseudoepitheliomatous hyperplasia. Papillary dermis revealed numerous ectatic lymphatics filled with eosinophilic proteinaceous material (Figures 3 and 4).



Fig. 1: Clinical image of vulva showing papulonodular lesion and sinus opening




Fig. 4: Numerous ectatic lymphatics lined by flattened endothelial cells in superficial dermis (H&E).



Fig. 2: Gross specimen of vulva showing pustules




Fig. 5: Sinus showing purulent exudative material (H&E)




Fig. 3: Papillary dermis showing numerous ectatic lymphatics filled with eosinophilic proteinaceous material (H&E)




Fig. 6: Chronic inflammation with multinucleated foreign body giant cell reaction

Areas of neutrophilic exocytosis and multiple sinuses lined by granulation tissue and containing neutrophilic microabscesses were seen (Figure 5). Reticular dermis and subcutaneous tissue showed diffuse extensive lymphoplasmacytic infiltration and multinucleated foreign body giant cell reaction (Figure 6). Focally, destruction of sebaceous and skin adnexal structures were also noted.

3. Discussion

Hidradenitis suppurativa (HS) is a chronic, inflammatory skin lesion, which tends to recur, may cause disfigurement, contractures and obstruction of superficial lymphatics vessels with collection of lymph in the interstitial space. Vulvar LC is known to develop as a complication of Hidradenitis suppurativa (HS).³ Lymphedema can develop in either a widespread or localized manner.⁴ It has been hypothesized due to recurrent, chronic inflammation and scarring, subcutaneous lymphatic channels are hindered from draining into the common lymphatic system completely. Thus leading to cumulation of lymph in the superficial interstitial tissue and furthermore smooth muscle around these lymphatic channels contract and causes vesicle formation.⁵

HS generally affects women between 30–58 years, with an average age of 46 years. However, acquired LC is frequently reported in slightly older women as compared to typical onset of HS, which is commonly reported in second and third decade of life. This is because of the fact that lymphatic obstruction occurs due to sequelae of longstanding HS.^{3,6}

Clinically, in HS patient presents with history of recurrent multiple pus discharging sinuses, scarring and semitranslucent skin-colored papules which coalescing into plaques, and bear a resemblance to gelatinous mass of ‘frog spawn’ on the surface of skin.^{1,5} It was traditionally known as HS polyposa.⁷

Radical surgery and /or radiation treatment for cervical cancer, infective diseases (acute cellulitis, sexually transmitted diseases, genital tuberculosis, erysiples, lymphogranuloma venereum), chronic inflammatory lesion; Crohn’s disease, bilateral varicose veins in the lower extremities, are the pathogenic factors for acquired LC.¹

Chu et al.⁴ employed the term “verrucous lymphostasis” to include both the generalized lymphedema of the abdomen and localized verrucous papules, plaques and nodules secondary to lymphedema on the buttocks, groin and labia majora in their five case series of HS.

Few cases of vulvar HS related acquired LC resembles human papilloma-related condyloma due to fluid filled vesicles along with concurrent lymphedema, which may expand to adjoining abdomen and thigh skin. Rarely, these lesions may have hyperkeratotic and warty appearance, which can mimic verrucous carcinoma.^{1,5} Various studies have shown that, there is a 50% increased chances of

malignancy patients with chronic HS. Especially, in HS patients the relative risk for squamous cell carcinoma is 4.6 and mostly confined to the perineal or buttock areas.^{3,8}

Histopathologic examination revealed thickened, hyperkeratotic overlying epidermis with elongated rete ridges. The superficial dermis displayed lymphangiectasia, which is characterized by multiloculated cystic spaces filled with lymphatic fluid and lined by endothelium.^{1,4}

Among 33 cases with genital lymphedema, Rosenweig et al.⁹ observed that most common cause of lymphatic obstruction to be metastatic malignancy of inguinal nodes, radiation therapy, and surgical excision of the lymph nodes. It is possible that HS is a rare cause of lymphedema.

The treatment of choice in patients with acquired vulvar lymphedema is excision of excess tissue followed by split-thickness skin grafting has been reported to be successful.¹

4. Conclusion

Vulva is the rare site for lymphangioma circumscriptum. It is important to consider LC among differential diagnoses of various papular/warty lesion of genital, in order to address the cosmetic and psychosexual implications and to offer adequate treatment.

5. Source of Funding

None.

6. Conflicts of interest

There are no conflicts of interest.

References

1. Kokcu A, Yildiz L, Bildircin D, Kandemir B. Vulvar lymphangioma circumscriptum presenting periodic symptoms. *BMJ*. 2010;201:620103056. doi:10.1136/bcr.06.2010.3056.
2. Moosbrugger EA, Mutasim DF. Hidradenitis suppurativa complicated by severe lymphedema and lymphangiectasias. *J Am Academy Dermatol*. 2011;64(6):1223–4.
3. Micieli R, Alavi A. Lymphedema in patients with hidradenitis suppurativa: a systematic review of published literature. *Int J Dermatol*. 2018;57(12):1471–80.
4. Chu EY, Kovarik CL, Lee RA. Lymphedematous verrucous changes simulating squamous cell carcinoma in long-standing hidradenitis suppurativa. *Int J Dermatol*. 2013;52(7):808–12.
5. Piernick DM, Mahmood SH, Daveluy S. Acquired lymphangioma circumscriptum of the genitals in an individual with chronic hidradenitis suppurativa. *J Am Acad Dermatol*. 2018;4(1):64–6.
6. Grada AA, Phillips TJ. Lymphedema: pathophysiology and clinical manifestations. *J Am Academy Dermatol*. 2017;77(6):1009–20.
7. Wrone DA, Landeck A, Dibbell DG, Xie H, Warner TF. Hidradenitis suppurativa polyposa. *Pathol Res Pract*. 2000;196(8):589–92. doi:10.1016/S0344-0338(00)80033-3.
8. Chapman S, Delgadillo DI, Barber C, Khachemoune A. Cutaneous squamous cell carcinoma complicating hidradenitis suppurativa: a review of the prevalence, pathogenesis, and treatment of this dreaded complication. *Acta Dermatovenerol Alp Pannonica Adriat*. 2018;27(1):25–8.
9. Rosenzweig LB, Brett AS, Lefavre JF, Vandersteenhoven JJ. Hidradenitis suppurativa complicated by squamous cell carcinoma

and paraneoplastic neuropathy. *Am J Med Sci.* 2005;329:150–2.
doi:10.1097/00000441-200503000-00008.

Mehak, Post Graduate Student

Sanjeev Uppal, Professor and HOD  <https://orcid.org/0000-0001-5200-9558>

Author biography

Neelam Gupta, Professor and HOD

Preeti Joseph, Associate Professor

Lav Behl, Post Graduate Student

Cite this article: Gupta N, Joseph P, Behl L, Mehak, Uppal S. Acquired lymphangioma circumscriptum of the vulva in a middle aged woman with chronic hidradenitis suppurativa – A case report. *IP J Diagn Pathol Oncol* 2023;8(1):46-49.